

Relic density in SO(10) SUSY GUT models

Katri Huitu* and Jari Laamanen†

**Department of Physics, and Helsinki Institute of Physics,
P.O. Box 64, FIN-00014 University of Helsinki, Finland*

†*Institut für Physik, Technische Universität Dortmund, D-44221 Dortmund, Germany*

Abstract. Non-universal boundary conditions in grand unified theories can lead to non-universal gaugino masses at the unification scale. In R -parity preserving theories the lightest supersymmetric particle is a natural candidate for the dark matter. In this talk the composition of the lightest neutralino is studied, when nonuniversal gaugino masses come from representations of SO(10). In these cases, the thermal relic density compatible with WMAP observations is found.

Keywords: Gaugino masses, relic density, dark matter

PACS: 12.60.Jv, 95.35.+d

INTRODUCTION

The phenomenology of supersymmetric models depends crucially on the compositions of neutralinos and charginos. In addition to the laboratory studies, relevant input is obtained from the dark matter searches, where the WMAP satellite has put precise limits on the relic density. Supersymmetric theories which preserve R -parity contain a natural candidate for the cold dark matter particle. If the lightest neutralino is the lightest supersymmetric particle (LSP), it can provide the appropriate relic density.

In many supergravity type models the lightest neutralino is bino-like, which often leads to too high thermal relic density, as compared to the limits provided by the WMAP experiment. When the gaugino masses are not universal at the grand unification scale, the resulting neutralino composition changes from the usual universal gaugino mass case [1].

In this talk, the thermal relic density of the neutralino LSP is studied, when gaugino masses are due to nonuniversal representations of SO(10) grand unified theory (GUT) [2, 3]. SO(10) has many attractive features when compared to SU(5), in which the relic density has been studied e.g. in [4]: the one family matter fermions can be put to one spinor representation of SO(10), including the right handed neutrino [5], representations are anomaly free, and R -parity may result from some gauge symmetry breaking, to mention some.

Gaugino masses originate from the non-renormalizable terms in the $N = 1$ supergravity Lagrangian involving the gauge kinetic function $f_{ab}(\Phi)$ [6]. The gauge part of the Lagrangian contains the gauge

kinetic function coupling with two field strength superfields W^a . The function $f_{ab}(\Phi)$ is an analytic function of the chiral superfields Φ in the theory. It transforms as a symmetric product of two adjoint representations under the GUT gauge group, since the field strength superfield transforms in the adjoint representation. In order to generate a mass term for the gauginos, the gauge kinetic function must be non-minimal, i.e. it must not be a constant [7]. The chiral superfields Φ consist of a set of gauge singlet superfields Φ^s and gauge nonsinglet superfields Φ^n , respectively, under the grand unified group. If the auxiliary part F_Φ of a chiral superfield Φ in the $f_{ab}(\Phi)$ gets a vev, then gaugino masses arise from the coupling of $f_{ab}(\Phi)$ with the field strength superfield W^a . The Lagrangian for the coupling of gauge kinetic function with the gauge field strength is written as

$$\mathcal{L}_{gk} = \int d^2\theta f_{ab}(\Phi) W^a W^b + \text{H.C.}, \quad (1)$$

where a and b are gauge group indices (for example, $a, b = 1, 2, \dots, 45$ for SO(10)), and repeated indices are summed over. The scalar fields in $f_{ab}(\Phi)$ are suppressed by the inverse powers of M_P , so the gauge kinetic function $f_{ab}(\Phi)$ can be expanded,

$$f_{ab}(\Phi) = f_0(\Phi^s) \delta_{ab} + \sum_n f_n(\Phi^s) \frac{\Phi^n}{M_P} + \dots, \quad (2)$$

where Φ^s and Φ^n are the singlet and nonsinglet chiral superfields, respectively. Here $f_0(\Phi^s)$ and $f_n(\Phi^s)$ are functions of gauge singlet superfields Φ^s , and M_P is some large scale. When F_Φ gets a vev $\langle F_\Phi \rangle$, the interaction (1) gives rise to gaugino masses:

$$\mathcal{L}_{gk} \supset \frac{\langle F_\Phi \rangle_{ab}}{M_P} \lambda^a \lambda^b + \text{H.C.}, \quad (3)$$

where $\lambda^{a,b}$ are gaugino fields. The non-universal gaugino masses are generated by the nonsinglet chiral superfield

e-mail: jari.laamanen@uni-dortmund.de

Φ^n that appears linearly in the gauge kinetic function $f_{ab}(\Phi)$ (2).

Gauginos belong to the adjoint representation of the gauge group, which in the case of $SO(10)$ is the **45** dimensional representation. Because Eq. (3) must be gauge invariant, Φ and F_Φ must belong to some of the following representations appearing in the symmetric product of the two **45** dimensional representations of $SO(10)$ [8]:

$$(\mathbf{45} \otimes \mathbf{45})_{\text{Sym}} = \mathbf{1} \oplus \mathbf{54} \oplus \mathbf{210} \oplus \mathbf{770}. \quad (4)$$

The representations **54**, **210** and **770** may lead to non-universal gaugino masses, while the **1** dimensional representation gives manifestly the universal gaugino masses. The relations between the gaugino masses are determined by the representation invariants, and are specific for each of the representations. Because the gauge kinetic function (2) can get contributions from several different Φ 's, a linear combination of any of the representations is also possible. In that case the gaugino masses are not anymore uniquely determined, in contrast to the contribution purely from one representation. Because of this it is more instructional to study the representations separately. In other words, we assume that the dominant component of the gaugino masses comes only from one representation, and more specifically, since we want to study non-universal gaugino masses, from one of the non-singlet representation.

DARK MATTER IN $SO(10)$ REPRESENTATIONS

Breaking Chains: $SO(10) \rightarrow H \rightarrow \text{SM}$

The GUT group $SO(10)$ breaks down to the Standard Model (SM) gauge group $SU(3) \times SU(2) \times U(1)$ via some intermediate gauge group H . Therefore the gaugino mass relations depend also on the gauge group breaking chain, in addition to the representation invariants coming from the gauge kinetic function. Moreover, the intermediate breaking scale affects also the generated gaugino masses. However, if the gauge breaking from the GUT scale to the SM scale takes place all at the GUT scale, these loop-induced messenger contributions [9] can be neglected in comparison to the tree-level contributions. Table 1 shows various possible $SO(10)$ breaking chains [10, 8] for the two chosen representations. Some of the subgroups lead to universal gaugino masses, or to massless gauginos. In this talk we suppose that the gauge breaking from $SO(10)$ to $SU(3) \times SU(2) \times U(1)$ happens at the GUT scale, and that the GUT breaking does not affect the gauge coupling unification. We also limit our study to the representations **54** and **210** and to three specific intermedi-

ate gauge groups, $SU(4) \times SU(2) \times SU(2)$, $SU(2) \times SO(7)$ and $SU(5) \times U(1)$. Table 2 displays the ratios of resulting

TABLE 1. Various breaking chains of $SO(10)$

F_Φ	H	Subgroup description
54	$SU(4) \times SU(2) \times SU(2)$	Pati–Salam
	$SU(2) \times SO(7)$	
	$SO(9)$	Universal gauginos
210	$SU(4) \times SU(2) \times SU(2)$	Massless gluino
	$SU(3) \times SU(2) \times SU(2) \times U(1)$	Massless $SU(2)_L$ gauginos
	$SU(3) \times SU(2) \times U(1) \times U(1)$	
	$SU(5) \times U(1)$	'Flipped' $SU(5)$

gaugino masses at tree level as they arise when F_Φ belongs to various representations of $SO(10)$. The relations at the electroweak scale resulting from 1-loop renormalization group running are also displayed.

TABLE 2. Ratios of the gaugino masses at the GUT scale in the normalization $M_3^{\text{GUT}} \equiv M_3(\text{GUT}) = 1$, and at the electroweak scale in the normalization $M_3^{\text{EW}} \equiv M_3(\text{EW}) = 1$

F_Φ	H	M_1^{GUT}	M_2^{GUT}	M_3^{GUT}	M_1^{EW}	M_2^{EW}	M_3^{EW}
1		1	1	1	0.14	0.29	1
54	$SU(4) \times SU(2) \times SU(2)$	-1	-1.5	1	-0.15	-0.44	1
54	$SU(2) \times SO(7)$	1	-7/3	1	0.15	-0.68	1
210	$SU(5) \times U(1)$	-96/25	1	1	-0.56	0.29	1

Representation 210

In the representation **210** we inspected the breaking chain through the intermediate gauge group $SU(5) \times U(1)$ (called flipped $SU(5)$). In Figure 1 the area of preferred thermal relic density in the representation **210** is plotted for a set of (GUT scale) parameters. For the chosen parameters, rather large WMAP preferred regions are found for the large values of M_2 and/or m_0 parameters. The dark shaded areas represent larger relic density than the lighter areas. wmap denoted filling is the WMAP preferred region, lep shows an area, where the experimental mass limits are not met, rge shows an area where there is no radiative EWSB, and lsp an area where neutralino is not the LSP. For the relic density, we use here the WMAP combined three year limits [11]

$$\Omega_{\text{CDM}} h^2 = 0.11054^{+0.00976}_{-0.00956} \quad (2\sigma). \quad (5)$$

The tiny area enclosed by the bsg contour is disallowed by $b \rightarrow s\gamma$ constraint. We have used the two sigma world average of $BR(b \rightarrow s\gamma) = (355 \pm 24^{+9}_{-10} \pm 3) \times 10^{-6}$ for the branching fraction [12]. The dash-dotted line (h) encloses an area with $m_h < 114$ GeV. The lep limits for the particle masses are the same as in [13]. The spectrum was calculated with SOFTSUSY [14] and relic densities and constraints with micrOMEGAS [15].




FIGURE 1. Relic density $\Omega_\chi h^2$ in the representation **210** and $H = \text{SU}(5) \times \text{U}(1)$ in (M_2, m_0) plane: $\tan\beta = 10$, $\text{sgn}(\mu) = +1$, $A_0 = 0$. The key is explained in the text.

The WMAP preferred region is very wide as compared to e.g. models in typical mSUGRA or $\text{SU}(5)$ [4, 13]. For a large part of the parameter space the relic density is below the WMAP limit. The lightest neutralino is either wino or higgsino, which explains the low overall relic density. In the diagonal of the figure the lightest chargino mass is very close to the lightest neutralino mass leading to the remarkable co-annihilation through the processes $\tilde{\chi}_1^0 \tilde{\chi}_1^\pm \rightarrow q_u \bar{q}_d$ and $\tilde{\chi}_1^0 \tilde{\chi}_1^0, \tilde{\chi}_1^\pm \tilde{\chi}_1^\pm \rightarrow q\bar{q}, \ell\bar{\ell}, W^+ W^-$. The spectrum is relatively heavy for the WMAP preferred region, but the value of μ stays around a TeV, which doesn't necessarily require large fine-tuning.

Representation 54

The two chains of the **54** dimensional representation do not lead to large areas with acceptable relic density. Because $\tilde{\chi}_1^0$ is mostly bino in $\text{SU}(2) \times \text{SO}(7)$ chain, the spectrum with preferred RD is quite light and conflicts with collider constraints in some parts of the parameter space (although the higgsino component keeps the overall thermal relic density lower than for example in the universal SUGRA model). In $\text{SU}(4) \times \text{SU}(2) \times \text{SU}(2)$ the higgsino component in the lightest neutralino is larger, and also co-annihilation with chargino is present at some points of parameter space, so the allowed region extends also to heavier spectrum. For more detailed analysis the reader is referred to an upcoming publication [16].

SUMMARY

We studied the dark matter allowed regions in the $\text{SO}(10)$ GUT representations, of which all but the singlet may lead to non-universal gaugino masses. The WMAP preferred relic density regions were quite distinct for different representations, thus leading to quite different particle spectra for each representation. As an example, we showed the behavior of the thermal relic density in the representation **210**, since there the spectrum is quite heavy as compared to the universal SUGRA case. It was also shown, that the WMAP preferred relic density area is very large in this case. Moreover, it is important to realize that there is no automatical theoretical preference for the gaugino masses to be unified.

ACKNOWLEDGMENTS

The work of K.H. is supported by the Academy of Finland (project No. 115032). The work of J.L. is supported by the Bundesministerium für Bildung und Forschung, Berlin-Bonn.

REFERENCES

1. K. Huitu, J. Laamanen, P. N. Pandita, and S. Roy, *Phys. Rev. D* **72**, 055013 (2005), [hep-ph/0502100](#).
2. H. Fritzsch, and P. Minkowski, *Ann. Phys.* **93**, 193–266 (1975).
3. Z.-Y. Zhao, *J. Phys.* **G8**, 1019 (1982).
4. K. Huitu, et al. (2008), [0808.3094](#).
5. R. N. Mohapatra, and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980).
6. E. Cremmer, S. Ferrara, L. Girardello, and A. Van Proeyen, *Phys. Lett.* **B116**, 231 (1982).
7. S. Ferrara, L. Girardello, and H. P. Nilles, *Phys. Lett.* **B125**, 457 (1983).
8. N. Chamoun, C.-S. Huang, C. Liu, and X.-H. Wu, *Nucl. Phys.* **B624**, 81–94 (2002), [hep-ph/0110332](#).
9. G. F. Giudice, and R. Rattazzi, *Nucl. Phys.* **B511**, 25–44 (1998), [hep-ph/9706540](#).
10. C. S. Aulakh, and R. N. Mohapatra, *Phys. Rev. D* **28**, 217 (1983).
11. D. N. Spergel, et al., *Astrophys. J. Suppl.* **170**, 377 (2007), [astro-ph/0603449](#).
12. E. Barberio, et al., Averages of b-hadron properties at the end of 2006 (2007), [0704.3575](#).
13. K. Huitu, J. Laamanen, and S. Roy (2007), [0710.2058](#).
14. B. C. Allanach, *Comput. Phys. Commun.* **143**, 305–331 (2002), [hep-ph/0104145](#).
15. G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput. Phys. Commun.* **176**, 367–382 (2007), [hep-ph/0607059](#).
16. K. Huitu, and J. Laamanen, Relic density in non-universal gaugino mass models with $\text{SO}(10)$ GUT symmetry (2008), in preparation.